

# Decay Results for Solutions to the Magneto-Hydrodynamics equations

M. E. Schonbek\*    T. P. Schonbek†    Endre Süli‡

January, 1995

**Key words:** Magneto-hydrodynamics equations, decay of solutions

**AMS(MOS) subject classification:** 35B40, 76W05

In this paper we describe the long-time behavior of the solutions to the viscous Magneto-Hydrodynamic equations. We will only mention the results obtained. The details of the proofs and corresponding references can be found in [3].

We study the Magneto-Hydrodynamic equations

$$\begin{aligned} \frac{\partial}{\partial t} u + (u \cdot \nabla) u - (B \cdot \nabla) B + \nabla p &= \Delta u + f, \\ \text{(MHD)} \quad \frac{\partial}{\partial t} B + (u \cdot \nabla) B - (B \cdot \nabla) u &= \Delta B, \\ \nabla \cdot u = 0, \quad \nabla \cdot B &= 0, \end{aligned}$$

supplemented with the initial conditions

$$u(x, 0) = u_0(x), \quad B(x, 0) = B_0(x),$$

for  $x \in \mathbf{R}^n$ ,  $2 \leq n \leq 4$ ,  $t \geq 0$ . We assume that the forcing function  $f$  is divergence free; i.e., that  $\nabla \cdot f(t) = 0$  for all  $t \geq 0$ . We show that solutions to the MHD equations, unlike solutions to the underlying heat equation, will

---

\*University of California, Santa Cruz, California. Research partially supported by NSF Grant No. DMS-9020941.

†Florida Atlantic University, Boca Raton, Florida.

‡Oxford University Computing Laboratory, 11 Keble Road, Oxford OX1 3QD, England.  
e-mail: endre.suli@comlab.ox.ac.uk

generically decay algebraically at a slow rate of  $(t + 1)^{-\alpha}$ , where  $\alpha = n/2$  or  $n/2 - 1$ , with  $n$  equal to the spatial dimension. The idea is that the non-linear terms will produce slow modes which will launch long waves preventing a too rapid decay. In other words, even if the data is highly oscillatory, the convection term will produce some mixing of the modes which will introduce long waves slowing down the decay.

More precisely, we show that weak solutions to the MHD equations, subject to large initial data outside a class of functions with total radially equidistributed energy, decay algebraically (rather than exponentially). In particular we prove that, for such solutions, the total energy (kinetic plus magnetic) and the magnetic energy have slowly decaying algebraic lower bounds. Moreover, we show in which cases the lower bounds are valid for the kinetic energy alone or the magnetic energy alone. Thus, our results reinforce mathematically the observation made by Chandrasekhar [1] that “the magnetic field in systems of large linear dimensions can endure for relatively long periods of time”.

We study upper and lower bounds. For the upper bounds we improve the results obtained by [2]. We show that

**THEOREM [A].** *Let the initial datum  $(u_0, B_0) \in H \times H$  and assume  $f \in L^1(0, \infty; L^2(\mathbf{R}^n)^n) \cap A_{n/4+1} \cap B_4 \cap C_{(n+3)/2}$ .*

**a)** *If  $n = 2$  and not all components of  $\hat{u}_0(0)$  or  $\hat{B}_0(0)$  are zero (in the sense defined at the beginning of Section 2), then*

$$\|u(t) - u_0(t)\|_2^2 + \|B(t) - B_0(t)\|_2^2 \leq C_{D_0}(t + 1)^{-n/2-1/2};$$

**b)** *if  $n \geq 3$ , or if  $n = 2$  and  $(u_0, B_0) \in [H \cap L^1(\mathbf{R}^n)^n]^2$ , then*

$$\|u(t) - u_0(t)\|_2^2 + \|B(t) - B_0(t)\|_2^2 \leq C_{D_1}(t + 1)^{-n/2-1}.$$

For the lower bounds we show

**THEOREM [B].** *Let  $(u_0, B_0) \in [W_2 \cap H]^2$ ,  $f \in L^1(0, \infty; L^2(\mathbf{R}^n)^n) \cap A_{n/4+1} \cap B_4 \cap C_{(n+3)/2}$ , and let  $(u(x, t), B(x, t))$  be a weak solution of the MHD equations with initial datum  $(u(x, 0), B(x, 0)) = (u_0(x), B_0(x))$ .*

**a)** *If  $\hat{u}_0(0) \neq 0$  or  $\hat{B}_0(0) \neq 0$ , then*

$$C_0(t + 1)^{-n/2} \leq \|u(\cdot, t)\|_2^2 + \|B(t)\|_2^2 \leq C_1(t + 1)^{-n/2}.$$

b) If  $(u_0, B_0) \in [W_2 \cap H \cap [L^1(\mathbf{R}^n)]^n]^2$  (so that

$$\hat{u}_0(0) = (2\pi)^{-n/2} \int_{\mathbf{R}^n} u_0(x) dx = 0, \hat{B}_0(0) = (2\pi)^{-n/2} \int_{\mathbf{R}^n} B_0(x) dx = 0,$$

by Borchers' Lemma) and  $(u, B) \notin \mathcal{M}_0$ , then

$$C_2(t+1)^{-n/2-1} \leq \|u(\cdot, t)\|_2^2 \leq C_3(t+1)^{-n/2-1}.$$

where Borchers' lemma was

LEMMA. Let  $u \in L^1(\mathbf{R}^n)^n \cap H$ . Then

$$\int_{\mathbf{R}^n} u dx = 0.$$

We have used the following notation .  $H^m(\mathbf{R}^n)$  denotes the Hilbertian Sobolev space on  $\mathbf{R}^n$  of index  $m$ ,  $m \geq 0$ ;  $L^p(\mathbf{R}^n)$  stands for the Lebesgue space equipped with its standard norm  $\|\cdot\|_p$ ,  $1 \leq p \leq \infty$ ;

$$\mathcal{V} = \{v \in [C_0^\infty(\mathbf{R}^n)]^n : \nabla \cdot v = 0\}, \quad H = \text{closure of } \mathcal{V} \text{ in } [L^2(\mathbf{R}^n)]^n.$$

$$W_1 = \{v : \int_{\mathbf{R}^n} |x|^2 |v(x)| dx < \infty\}, \quad W_2 = \{v : \int_{\mathbf{R}^n} |x| |v(x)|^2 dx < \infty\},$$

Moreover, if  $\mu, \nu, \sigma \in \mathbf{R}$ , we say

**a**  $f \in A_\mu$  if there exists  $C \geq 0$  such that

$$\|f(t)\|_2 \leq C(t+1)^{-\mu} \quad \text{for } t \geq 0.$$

**b**  $f \in B_\sigma$  if there exists  $C \geq 0$  such that

$$|\hat{f}(\xi, t)| \leq C|\xi|^\sigma \quad \text{for } t \geq 0, \xi \in \mathbf{R}^n.$$

**c**  $f \in C_\nu$  if there exists  $C \geq 0$  such that

$$\|f(t)\|_\infty \leq C(t+1)^{-\nu} \quad \text{for } t \geq 0.$$

Finally, Given  $u = (u_1, \dots, u_n)$  and  $B = (B_1, \dots, B_n)$  in  $[L^1(0, \infty; L^2(\mathbf{R}^n))]^n$ , let

$$\tilde{\mathcal{A}}_{ij} = \int_0^\infty \int_{\mathbf{R}^n} (u_i u_j - B_i B_j) dx,$$

$$\tilde{\mathcal{C}}_{ij} = \int_0^\infty \int_{\mathbf{R}^n} (u_i B_j - B_i u_j) dx.$$

Then, introducing the matrices  $\tilde{\mathcal{A}} = [\tilde{\mathcal{A}}_{ij}]$  and  $\tilde{\mathcal{C}} = [\tilde{\mathcal{C}}_{ij}]$ , we define

$$\mathcal{M}_0 = \{(u, B) \in [L^2(\mathbf{R}^n)]^{2n} : \tilde{\mathcal{A}} \text{ is scalar and } \tilde{\mathcal{C}} = 0\}.$$

The idea of the proof is to compare the decay of the solutions to MHD with the decay of the solution to the underlying heat system. It is first shown that solutions to the heat equation such that the Fourier transform of the initial data has a zero of order  $k$  at the origin, decay at most like  $(t+1)^{-k/2-n/4}$ ; that is, if

$$|\hat{u}_0(\xi)| = C_0 |\xi|^k + O(|\xi|^{k+1}),$$

then the solution  $v$  of  $v_t = \Delta v$ ,  $v(x, 0) = u_0(x)$  satisfies

$$(1) \quad C_0(t+1)^{-n/2-k} \leq \|v(\cdot, t)\|_2^2 \leq C_0(t+1)^{-n/2-k}$$

for  $t \geq 0$ . By an argument similar to Wiegner's for solutions to the Navier-Stokes equations, it is easy to show that if  $V$  satisfies  $V_t = \Delta V$ ,  $V(x, 0) = (u_0(x), B_0(x))$  and  $W = V - (u, B)$ , then

$$(2) \quad \|W(t)\|_2^2 \leq C_W(t+1)^{-n/2-\beta}$$

where  $\beta = 1/2$  or  $\beta = 1$  depending on whether  $n = 2$  or  $n \geq 3$ , respectively. Thus, for the case in which  $\int_{\mathbf{R}^n} u_0 dx \neq 0$ ; i.e.,  $\hat{u}_0(0) \neq 0$ , it is immediate from (1) and (2) that

$$\|u(t)\|_2^2 + \|B(t)\|_2^2 \geq C(t+1)^{-n/2}.$$

When the Fourier transform of the initial data has a zero of higher order the conditions from the theorem are such that a long wave is always present. That is, we can compare the solution to the MHD equations with data at a large time  $T$  with the corresponding solution  $V_T$  to the heat system with data at time  $T$ . In this case, because of the long wave, it follows that

$$\|V_T(\cdot, t)\|_2^2 \geq C_T(t+1)^{-n/2-1}.$$

Setting  $W_T(\cdot, t) = V_T(\cdot, t) - (u(\cdot, t+T), B(\cdot, t+T))$ , we'll have our lower bounds if we can show that  $C_T \geq C_{W_T}$ . Since  $C_{W_T}$  will depend on the  $L^2$  norm of the data  $(u(T), B(T))$ ,  $T$  sufficiently large, it will follow that  $C_{W_T}$  is as small as needed. Combining once more the lower bound of  $V_T$  with the upper bound of  $W_T$ , we get

$$\|u(t)\|_2^2 + \|B(t)\|_2^2 \geq C(t+1)^{-n/2-1}.$$

## References

- [1] S. CHANDRASEKHAR, *Hydrodynamic and Hydromagnetic Stability*, Clarendon Press, Oxford, 1961.
- [2] S.D. MOHGOONER AND R.E. SARAYKER,  $L^2$  decay for solutions to the *MHD equations*, Journal of Mathematical and Physical Sciences, **23**, No 1, (1989), 35–53.
- [3] Decay results for Solutions to the Magneto Hydrodynamics equations.(with T.P. Schonbek and E. Suli) *Mathematical Analysis of Phenomena in Fluids and Plasma Dynamics*. Editors: K. Asano, A. Tani, S. Ukai. RIMS Kokyuroku 914, 98-102, (1995) .